

source: <https://doi.org/10.7892/boris.53271> | downloaded: 5.5.2023

1 **Taxon-specific $\delta^{13}\text{C}$ analysis of chitinous invertebrate remains in sediments from**
2 **Strandsjön, Sweden**
3
4
5
6
7 4 M. van Hardenbroek^{1,2}, A.F. Lotter², D. Bastviken³, T.J. Andersen⁴, O. Heiri¹
8
9
10 5
11
12 6 ¹ Institute of Plant Sciences and Oeschger Centre for Climate Change Research,
13
14 7 University of Bern, Altenbergrain 21, CH-3013 Bern, Switzerland.
15
16 8 ² Palaeoecology, Department of Physical Geography, Laboratory of Palaeobotany and
17
18 9 Palynology, Utrecht University, Budapestlaan 4, 3584 CD Utrecht, The Netherlands.
19
20 10 ³ Department of Thematic Studies - Water and Environmental Studies, Linköping
21
22 11 University, 58183 Linköping, Sweden.
23
24 12 ⁴ Department of Geography and Geology, University of Copenhagen, DK-1350
25
26 13 Copenhagen K, Denmark.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

15 **Abstract**
16
17 Taxon-specific stable carbon isotope ($\delta^{13}\text{C}$) analysis of chitinous remains of
18 invertebrates can provide valuable information about the carbon sources used by
19 invertebrates living in specific habitats of lake ecosystems (for example, sediments,
20 water column, or aquatic vegetation). This is complementary to $\delta^{13}\text{C}$ of sedimentary
21 organic matter (SOM), which provides an integrated signal of organic matter
22 produced in a lake and its catchment, and of diagenetic processes within sediments. In
23 a sediment record from Strandsjön (Sweden) covering the past circa 140 years, we
24 analyzed SOM geochemistry ($\delta^{13}\text{C}$, $\text{C:N}_{\text{atomic}}$, organic carbon content) and $\delta^{13}\text{C}$ of
25 chitinous invertebrate remains in order to examine whether taxon-specific $\delta^{13}\text{C}$

records could be developed for different invertebrate groups and whether these analyses provide insights into past changes of organic carbon sources for lacustrine invertebrates available in benthic and planktonic compartments of the lake. Invertebrate taxa included benthic chironomids (*Chironomus*, Chironomini excluding *Chironomus*, Tanytarsini, and Tanypodinae), filter-feeders on suspended particulate organic matter (*Daphnia*, *Plumatella* and *Cristatella mucedo*), and Rhabdocoela. $\delta^{13}\text{C}$ of chironomid remains indicated periodic availability of ^{13}C -depleted carbon sources in the benthic environment of the lake as $\delta^{13}\text{C}$ values of the different chironomid taxa fluctuated simultaneously between -34.7 and -30.5‰ (VPDB). *Daphnia* and Bryozoa showed parallel changes in their $\delta^{13}\text{C}$ values which did not coincide with variations in $\delta^{13}\text{C}$ of chironomids, though, and a 2-3‰ decrease since circa AD 1960. The decrease in $\delta^{13}\text{C}$ of *Daphnia* and Bryozoa could indicate a decrease in phytoplankton $\delta^{13}\text{C}$ as a result of lower lake productivity, which is in accordance with historical information about the lake that suggests a shift to less eutrophic conditions after AD 1960. In contrast, Rhabdocoela cocoons were characterized by relatively high $\delta^{13}\text{C}$ values (-30.4 to -28.2‰) that did not show a strong temporal trend, which could be related to the predatory feeding mode and wide prey spectrum of this organism group. The taxon-specific $\delta^{13}\text{C}$ analyses of invertebrate remains indicated that different carbon sources were available for the benthic chironomid larvae than for the filter-feeding *Daphnia* and bryozoans. Our results therefore demonstrate that taxon-specific analysis of $\delta^{13}\text{C}$ of organic invertebrate remains can provide complementary information to measurements on bulk SOM and that $\delta^{13}\text{C}$ of invertebrate remains may allow the reconstruction of past changes in carbon sources and their $\delta^{13}\text{C}$ in different habitats of lake ecosystems.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Keywords: invertebrates, chitinous remains, lake sediment, stable carbon isotopes,
sedimentary organic matter

Introduction

The stable carbon isotope composition of bulk organic matter in lake sediments is determined by the $\delta^{13}\text{C}$ values of different organic matter sources available in a lake and its catchment and by the preservation of the various inputs (Meyers and Teranes 2001). Environmental changes can affect the isotopic composition of organic matter, the amount of organic matter produced within lakes, the relative contribution of autochthonous and allochthonous organic matter, and the transport processes leading to the incorporation of organic matter into lake sediments. Therefore, analysis of $\delta^{13}\text{C}$ of bulk sedimentary organic matter (SOM) can potentially provide information on past changes in climate, productivity, origin of organic matter, pollution, and in-lake carbon cycling (Leavitt et al. 2006; Leng et al. 2005; Meyers and Lallier-Vergès 1999). It is difficult, however, to reconstruct limnological changes in different compartments (benthic, pelagic) of a lake based on $\delta^{13}\text{C}$ of SOM only, since variations in $\delta^{13}\text{C}$ of SOM may reflect variations in the isotopic composition of organic matter within the lake as well as in the lake catchment. Even if the approach is applied to lakes in which SOM originates mainly from lacustrine sources, variations in bulk SOM isotopic composition may reflect changes in carbon sources and organic matter production in the littoral, pelagial or profundal of the lake and selective preservation/degradation of various compounds and tissues. A possible approach to circumvent this problem is to analyze organic remains of selected organisms

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

76 originating from the lake itself. If these remains can be identified and associated with
77 organism groups with clear habitat preferences it may be possible to determine past
78 changes in the carbon isotopic composition of aquatic organic matter in specific
79 habitats and compartments of lakes.

80 Recently, a number of studies analyzed $\delta^{13}\text{C}$ of specific organic remains that
81 preserve well in lake sediments. These include remains of primary producers such as
82 *Potamogeton* (Herzschuh et al. 2010; Turney 1999) and traces of organic matter
83 within diatom valves (Barker et al. 2013; Hurrell et al. 2011) as well as of secondary
84 producers such as aquatic invertebrates (for example, cladocerans (Perga 2010) and
85 chironomids (Wooller et al. 2008)). Chitinous invertebrate remains largely consist of
86 proteins and chitin and are chemically robust and relatively resistant to microbial
87 degradation, especially if buried under anoxic conditions (Verbruggen et al. 2010).
88 They can be identified and associated with a particular group of organisms. Therefore,
89 it can be ensured that the analysed remains originate from the lake itself. Since
90 different invertebrate groups are characterized by different feeding modes (for
91 example, filter-feeders, deposit-feeders, or predators) and habitat preferences
92 (planktonic or benthic, littoral or profundal), taxon-specific analyses of invertebrate
93 $\delta^{13}\text{C}$ can potentially provide insights on processes that are active in different
94 compartments of lakes and affect the lacustrine food web and carbon cycle.

95 Until very recently chitinous aquatic invertebrate remains have received little
96 attention with regard to carbon isotope analysis. Perga (2010) demonstrated that
97 strong correlations exist between $\delta^{13}\text{C}$ values of tissue of living cladocerans of the
98 genus *Bosmina* and *Daphnia* and their chitinous carapaces. However, she concluded
99 that taphonomic processes should nevertheless be taken into account when
100 interpreting palaeoecological records as they can cause lower $\delta^{13}\text{C}$ (<1‰) of *Daphnia*

remains, because isotopically light C compounds seem to be selectively removed by microbial degradation (Perga 2011). Wooller et al. (2008) were the first to analyze $\delta^{13}\text{C}$ of head capsules of chironomid larvae in a sediment sequence from an Icelandic lake. They reported similar patterns in stable isotope values of chironomids and SOM. Van Hardenbroek et al. (2010a) studied the effects of various chemical pre-treatments commonly used in palaeolimnological studies on carbon isotopic measurements on chitinous remains of chironomid larvae and showed that these treatments had little effect on the $\delta^{13}\text{C}$ of the larval head capsules. These authors also demonstrated that head capsule $\delta^{13}\text{C}$ is influenced by the chironomid diet and that methanogenic carbon can influence the carbon isotopic signature of chironomid exoskeletons. Heiri et al. (2012) and Frossard et al. (2013) demonstrated that $\delta^{13}\text{C}$ values of chironomid head capsules are closely related to the carbon isotopic composition of the larval body and the food ingested by chironomid larvae.

Only very few studies are available that examine $\delta^{13}\text{C}$ of remains of multiple invertebrate groups in the same sediment samples. Van Hardenbroek et al. (2012; 2013) examined the relationship of $\delta^{13}\text{C}$ values of remains of five chironomid taxa and the resting eggs (ephippia) of water fleas of the genus *Daphnia* with methane fluxes in 17 lakes in Sweden and Siberia and reported a statistically significant relationship between diffusive methane flux measured at the lake surface and $\delta^{13}\text{C}$ of the fossils of some chironomid groups and of *Daphnia* ephippia. Wooller et al. (2012) and van Hardenbroek et al. (2013) presented down-core records of chironomid and *Daphnia* $\delta^{13}\text{C}$ from thermokarst lakes in Alaska and Siberia, respectively, which suggest major changes in the carbon cycle and methane availability in these lakes. Similar results were obtained by Frossard et al. (2014) using sediment cores from Lake Annecy (France) in which they observed decreasing $\delta^{13}\text{C}$ values in remains of

126 pelagic Cladocera and three chironomid taxa during 1930s -1950s. Frossard et al.
127 (2014) suggest that eutrophication and increased hypoxia led to increased
128 incorporation of ^{13}C -depleted carbon (microbially respired and/or methane-derived)
129 into the invertebrate tissues, which explains the lower $\delta^{13}\text{C}$ values in the invertebrate
130 remains.

131 Down-core studies examining $\delta^{13}\text{C}$ in the remains of multiple planktonic and
132 benthic invertebrate groups such as cladocerans, bryozoans, chironomids, and other
133 aquatic insects allows the taxon-specific records to be compared among indicator
134 groups in order to detect variations in $\delta^{13}\text{C}$ common to several groups. Hence, past
135 changes in the carbon cycle of lakes tracked by different invertebrates originating
136 from the benthic or pelagic part of lakes could be separated from background
137 variations in $\delta^{13}\text{C}$ common to all indicators. Here we present a first study presenting
138 $\delta^{13}\text{C}$ records based not only on the chitinous remains of Chironomidae (non-biting
139 midges) and Cladocera (water fleas), but also on the remains of Bryozoa (moss
140 animals), and Rhabdocoela (Turbellaria: free-living flatworms). Analyses were made
141 on sediments collected from Strandsjön, a small shallow lake in South-central Sweden
142 ($59^{\circ}52'28''$ N and $17^{\circ}10'5''$ E, 51 m a.s.l.). The catchment contains pine forests and
143 pastures, some with dairy farming since the beginning of the 20th century that led to
144 increased nutrient loading of the lake until the installation of waste water treatment in
145 the 1960s (Brunberg and Blomqvist 1998) and TP values were lowered to $41.3 \mu\text{g L}^{-1}$
146 in 2008 (van Hardenbroek et al. 2012). The history of eutrophication and recovery of
147 Strandsjön was expected to have affected $\delta^{13}\text{C}$ values of algae, as these are usually
148 ^{13}C -enriched when productivity is higher (Brenner et al. 1993; Hollander et al., 1993).
149 We therefore expected productivity-related changes in algal $\delta^{13}\text{C}$ to be reflected in
150 $\delta^{13}\text{C}$ of algivorous filter-feeders like *Daphnia* and Bryozoa and their remains. In

contrast, we expected that $\delta^{13}\text{C}$ values would be not (or less clearly) related to past changes in productivity for predatory Rhabdocoela or for benthic chironomids, the latter feeding on a mixture of detritus from the water column and decomposed SOM. Strandsjön was selected because $\delta^{13}\text{C}$ of invertebrate remains in surface sediments from this lake were previously analysed by van Hardenbroek et al. (2012), providing additional information for the interpretation of invertebrate $\delta^{13}\text{C}$ in this down core study.

Materials and Methods

Sediment characteristics

The coring location was chosen at 2.4 m water depth (Fig. 1). This location represents the deepest part of a transect of surface sediment samples obtained from the lake in which $\delta^{13}\text{C}$ of invertebrate remains were analyzed (van Hardenbroek et al. 2012). A 36 cm long core with undisturbed sediment-water interface was obtained using a gravity corer (Uwitec, Austria) and sub-sampled in the field at 1 cm resolution. The samples were stored in plastic bags and kept cool and dark until freeze-drying upon arrival in the laboratory. Freeze-dried samples were used for dating by gamma spectrometry using a Canberra low-background Ge-well detector. ^{210}Pb was measured via its gamma-peak at 46.5 keV, ^{226}Ra via the granddaughter ^{214}Pb (peaks at 295 and 352 keV), and ^{137}Cs via its peak at 661 keV.

Samples for stable isotope analysis of SOM were rinsed in 2.5% HCl for 15 minutes to remove carbonates, rinsed three times with demineralized water,

centrifuged 4 min at 2000 rpm to remove excess water, and freeze-dried. C:N_{atomic} ratios, as well as stable carbon isotopes of SOM were analyzed on a PDZ Europa ANCA-GSL elemental analyzer coupled to a PDZ Europa 20-20 IRMS. Two laboratory standards (nylon and peach leaves) of known relation to international standards for VPDB were used as references. Replicate sample measurements on standards (n = 43) indicated an analytical error (2σ) of ±0.05‰ for δ¹³C.

δ¹³C of invertebrate remains

Samples for δ¹³C analysis of chitinous invertebrate remains were deflocculated in 10% KOH for 2 h at room temperature and sieved with 200- and 100-μm sieves (van Hardenbroek et al. 2010b). Rinsing with an alkaline solution has the added advantage of removing adsorbed fulvic and humic acids (Stevenson 1982). Sieve residues were soaked in 2.5% HCl for 15 minutes, rinsed three times and stored in demineralized water in the dark. Remains were identified under a dissecting microscope at 40–100x magnification following Wood and Okamura (2005) for Bryozoa statoblasts, Vanderkerkhove et al. (2004) for the resting stages of *Daphnia*, Brooks et al. (2007) for chironomid head capsules, and Luther (1955) and Frey (1964) for Rhabdocoela cocoons. Chironomid head capsules were separated into the categories *Chironomus* spp., Chironomini (excluding *Chironomus*), Tanytarsini, Orthoclaadiinae, and Tanypodinae. Furthermore, we differentiated ephippia of the cladoceran genus *Daphnia*, statoblasts of the Bryozoa taxa *Plumatella* and *Cristatella mucedo*, and the chitinous cocoons of Rhabdocoela.

After identification, remains were transferred with forceps directly into pre-weighed ultraclean tin cups. Tin cups were dried on a hotplate at 50°C for 24 h after

which they were re-weighed and crimped for stable isotope analysis. The average number of remains in a sample was 18 for *Chironomus* (average sample weight 23 µg), 36 for other Chironomini (33 µg), 45 for Tanytarsini (24 µg), 22 for Tanypodinae (22 µg), 34 for *Daphnia* (40 µg), 41 for *Plumatella* (18 µg), 1.2 for *C. mucedo* (28 µg), and 40 for Rhabocoela (47 µg) (Supplementary Table S1). Control samples of water from sieve residues were evaporated in tin cups and no carbon contamination was detected. Samples of invertebrate remains were analyzed on a Fisons NA 1500 NCS Elemental Analyzer coupled to a Thermo Electron Delta plus IRMS. A laboratory standard (Naxos GQ graphite quartzite) of known relation to international standards for VPDB was used as reference. Replicate sample measurements of the secondary standard (n = 30) indicated an analytical error (2σ) of ±0.05%.

Results

Age-depth model

High ¹³⁷Cs-activity (>2000 Bq kg⁻¹) was measured above 4.5 cm sediment depth in the Strandsjön sediments pointing to a link to the 1986 Chernobyl accident for this material. This suggests that the sediment at 4.5 cm depth was deposited in 1986 or shortly thereafter. ¹³⁷Cs-profiles measured in European lake sediments typically feature a second maximum at AD 1963, coinciding with the peak in above-ground nuclear bomb testing (Appleby 2001), but in Strandsjön this feature seems to be hidden by slight downward penetration of the high ¹³⁷Cs-activity in the top 4.5 cm. We measured surface contents of unsupported ²¹⁰Pb of approximately 220 Bq kg⁻¹. ²¹⁰Pb contents decreased more or less exponentially down core with the exception of

two outliers at 17-18 and 25-26 cm depth. The calculated flux of unsupported ^{210}Pb is $62 \text{ Bq m}^{-2} \text{ y}^{-1}$, which is similar to the supply rate of Knud Sø in Denmark ($58 \text{ Bq m}^{-2} \text{ y}^{-1}$) presented by Appleby (2001). Age-depth modeling using ^{210}Pb was based on a modified constant rate of supply (CRS) model (Appleby and Oldfield 1978) with 4.5 cm fixed at 1986 as a reference point (Fig. 2). The inventory below 25 cm was calculated on the basis of a regression of the content of unsupported ^{210}Pb versus accumulated mass above this level. The age-model was extrapolated for samples below 31 cm, resulting in an estimated age for the lowermost sample of circa AD 1870 ± 15 years. Due to the outliers in the ^{210}Pb measurements, we consider the CRS model less reliable in the deeper section of the core, which is why we report approximate ages for the record.

Sediment geochemistry and $\delta^{13}\text{C}$ of invertebrate remains

SOM $\delta^{13}\text{C}$ decreased from approximately -30 to -33‰ throughout the core and the C:N_{atomic} ratio showed a small, gradual decrease from approximately 11 to 9 (Fig. 3). The carbon content of the sediment was 15-20 % by weight (wt.%) in the lower part of the record up to 15 cm depth, then decreased to 10.5 wt.% at 10 cm depth, and increased again to 12.6 wt.% in the top 7 cm.

$\delta^{13}\text{C}$ values of filter-feeders that retrieve their food directly from the water column (*Daphnia*, *C. mucedo*, and *Plumatella*) remained relatively stable for much of the record (approximately 12-35 cm depth). Their $\delta^{13}\text{C}$ then decreased by 2-3‰ between approximately 5 and 10 cm depth (AD ~1960 to 1980), and increased again by approximately 1-1.5‰ in the uppermost sediments. *C. mucedo* had lowest $\delta^{13}\text{C}$ values (-35.1 to -31.2‰), followed by *Daphnia* (-32.8 to -30.5‰) and *Plumatella* (-

33.1 to -28.0‰) (Fig. 4). $\delta^{13}\text{C}$ of larval remains of chironomids showed variations at a higher frequency ranging between -34.6‰ and -30.3‰ (Fig. 3). Lowest values were recorded by *Chironomus* (-34.6 to -30.8‰), followed by Tanytarsini (-33.9 to 30.3‰), Tanypodinae (-33.8 to -31.0‰) and Chironomini other than *Chironomus* (-33.5 to -30.5‰) (Fig. 4). For most chironomid groups, relative shifts in $\delta^{13}\text{C}$ values were very similar with local minima in $\delta^{13}\text{C}$ recorded at 27.5, 20.5, 12.5, and 2.5-4.5 cm sediment depth (AD ~1890, ~1920, ~1950, and ~1990, respectively). The exception are the remains of *Chironomus*, which are present in the sediments between 0-19 cm and at 32 cm depth, as they show higher amplitude variations than the other chironomid groups. $\delta^{13}\text{C}$ of Rhabdocoela cocoons showed less temporal variability and was around -29.5‰ (-30.2 to -28.4‰) throughout the core.

Discussion

Feeding ecology of invertebrate groups

For the interpretation of $\delta^{13}\text{C}$ values it is important to understand the feeding ecology and habitat of the invertebrates in this study. The larvae of Chironomidae are benthic and, in lakes, live in or on sediments, or on hard substrates such as rocks, submerged wood and aquatic macrophytes. The Tanytarsini are typically algivores and deposit-feeders with limited capabilities for burrowing into sediments (Merritt et al. 2008; Mihuc and Toetz 1994; Vander Zanden and Rasmussen 1999). The Chironomini include a number of taxa that tend to burrow deeper into sediments, or in some cases into aquatic macrophytes. They feed predominantly as deposit-feeders, although some groups also filter particles from water which is pumped through the tubes the larvae

live in (Merritt et al. 2008; Moller Pillot 2009; Moog 2002), a habitat where they can also access ^{13}C -depleted biomass of methane oxidizing bacteria (Jones et al. 2008). The Tanypodinae are mainly free-living and mobile predators. However, several Tanypodinae are known to also feed on the substrate the larvae live on (Merritt et al. 2008; Vallenduuk and Moller Pillot 2007). Bryozoa live predominantly as colonies attached to hard substrates and feed on suspended nanoplanktonic algae from the water column, but they cannot move through the water themselves (Kaminski 1984; Okamura and Hatton-Ellis 1995; Wood and Okamura 2005). *Daphnia* are mobile zooplankton feeding on fine particles in the water column that range between 0.5 to 30 μm in diameter and include phytoplankton and bacteria (Geller & Müller 1981). It can be assumed, therefore, that $\delta^{13}\text{C}$ of *Daphnia* and Bryozoa reflects the material they filter from the water column. Rhabdocoela are active predators feeding on small invertebrates and large ciliate protozoans as well as scavengers (Jennings 1957; Kolasa and Tyler 2010). In contrast to benthic chironomids and filter-feeding *Daphnia* and bryozoa, rhabdocoels are probably less directly affected by changes in $\delta^{13}\text{C}$ values of phytoplankton or SOM but have a $\delta^{13}\text{C}$ signature representing a mixture of their prey organisms.

Differences in $\delta^{13}\text{C}$ offsets between invertebrate taxa

The carbon isotopic composition of invertebrate remains in Strandsjön showed some variability between the different invertebrate groups (Fig. 4). Similar to the findings of van Hardenbroek et al. (2013), the remains of *Chironomus* were ^{13}C -depleted relative to other chironomids and invertebrate groups. In Strandsjön, $\delta^{13}\text{C}$ values of *Chironomus* head capsules were only slightly more negative than the remains of other

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

301 Chironomidae and of *Daphnia*, which in turn tended to be only slightly more negative
302 than the $\delta^{13}\text{C}$ values recorded for bulk SOM. Interestingly, $\delta^{13}\text{C}$ values recorded for
303 statoblasts of the bryozoan *C. mucedo* were more negative than those of chironomid
304 remains in most samples, whereas the remains of *Plumatella* and of Rhabdocoela
305 cocoons were distinctly ^{13}C -enriched relatively to SOM and other invertebrates. We
306 found no published $\delta^{13}\text{C}$ values of sedimentary remains of *Plumatella* and
307 Rhabdocoela and only one published study with few $\delta^{13}\text{C}$ measurements on *C.*
308 *mucedo* from two sediment records in the United Kingdom (Turney, 1999). In this
309 study four $\delta^{13}\text{C}$ values of *C. mucedo* statoblasts from Llanilid are presented, ranging
310 between -29 and -25‰, approximately 0-1‰ lower than $\delta^{13}\text{C}$ values of SOM. For the
311 second site, Gransmoor, two $\delta^{13}\text{C}$ values of *C. mucedo* statoblasts are between -24
312 and -22‰ and approximately 0-4 ‰ higher than $\delta^{13}\text{C}$ of SOM. The variability in $\delta^{13}\text{C}$
313 offset between *C. mucedo* and SOM and the lack of stable isotope data clearly
314 indicates more work is required to understand which carbon sources are used by
315 Bryozoa and Rhabdocoela before their $\delta^{13}\text{C}$ values can be interpreted conclusively.

316 We expected that $\delta^{13}\text{C}$ offsets between taxa largely reflect differences in stable
317 isotope composition of their food sources. It must be considered, however, that
318 biochemical processes during the synthesis of the body tissues may also potentially
319 influence invertebrate $\delta^{13}\text{C}$ and can play a role when comparing remains from
320 different invertebrate taxa. Generally, the $\delta^{13}\text{C}$ of chitinous invertebrate structures is
321 closely related to the carbon isotopic composition of the soft tissue of the
322 invertebrates in question and of the food available for the animals (DeNiro and
323 Epstein 1978; Rau 1980). Schimmelmann (2011) reviewed the isotopic difference for
324 eight invertebrate-diet combinations and found that the isotopic difference between
325 diet and the corresponding D-glucosamine hydrochloride from invertebrate chitin

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

326 ranges between -2.2 and $+2.1\text{‰}$ (average $0.0 \pm 1.5\text{‰}$). His results suggest that taxa
327 can differ in the offset between $\delta^{13}\text{C}$ of ingested carbon and $\delta^{13}\text{C}$ of their chitinous
328 structures, but that this variation is mostly limited to approximately 0-2‰.
329 Experiments with laboratory cultures of *Chironomus riparius* indicated a ^{13}C -
330 depletion of 4th instar larval head capsules compared with 4th instar larval body tissue
331 of 1-2‰ (Heiri et al. 2012) and $0.9 \pm 0.2\text{‰}$ (Frossard et al. 2013). Perga (2011)
332 demonstrated that the offset in $\delta^{13}\text{C}$ between whole *Daphnia* and their ephippia
333 collected from the same water sample was within the standard deviation (0.2‰) of
334 replicate $\delta^{13}\text{C}$ measurements. Similarly, Macko et al. (1989) reported a 0.4-0.8‰ ^{13}C -
335 depletion of carapaces compared to body tissue of three marine crustaceans. As
336 mentioned before, little to nothing has been published about the stable isotope
337 composition of Bryozoa and Rhabdocoela or their remains. The large difference in
338 $\delta^{13}\text{C}$ values of remains of Rhabdocoela and *Plumatella*, relative to other invertebrate
339 remains examined in our record suggests that these taxa were able to access different
340 carbon sources than the other analysed invertebrate taxa. This interpretation is
341 supported by the observation that an offset is apparent even between *Plumatella* and
342 *C. mucedo*, both bryozoans, for which a similar offset between food, body tissue and
343 statoblasts can be expected. In our record *Plumatella* had on average 3.5‰ higher
344 $\delta^{13}\text{C}$ than *C. mucedo* and it is possible that this was the result of a difference in the
345 $\delta^{13}\text{C}$ of the particles ingested by these two taxa when filtering the water. Kaminski
346 (1984) demonstrates that *C. mucedo* selects small seston ($<7\text{ }\mu\text{m}$ in diameter), which
347 can include bacteria, whereas *Plumatella repens* prefers slightly larger particles
348 (ranging from 5 to 17 μm in diameter).
349
350 Trends in $\delta^{13}\text{C}$ of invertebrate remains

351

352 The four chironomid taxa showed similar variations in $\delta^{13}\text{C}$ with several minor
353 minima during the past circa 140 years (Fig. 3). All analyzed chironomid groups
354 featured the same type of high frequency signal, which makes it very unlikely that
355 these variations in $\delta^{13}\text{C}$ are related to changes in taxonomic composition that
356 happened simultaneously within all four groups. Since *Chironomus*, other
357 Chironomini, and Tanytarsini are largely detritivores and partly grazers and filterers,
358 and Tanypodinae are predators, it can be expected that the $\delta^{13}\text{C}$ of the chironomid
359 remains analyzed here mostly reflected the $\delta^{13}\text{C}$ of sedimentary detritus and
360 detritivorous prey animals. $\delta^{13}\text{C}$ of SOM in the Strandsjön record decreased gradually
361 between 35 and 0 cm depth. However, chironomids did not show the same decreasing
362 trend in $\delta^{13}\text{C}$ that we observed for SOM. This suggests that the invertebrates were
363 selectively feeding on and digesting certain organic matter components in the
364 sediments. Wooller et al. (2008; 2012) observed trends in $\delta^{13}\text{C}$ of chironomid remains
365 that were roughly similar to the patterns observed for SOM. In these studies, however,
366 Wooller et al. (2008; 2012) did not include taxonomic information about the analyzed
367 chironomids. This makes a direct comparison between their records and our study
368 difficult, since not all chironomid tribes and subfamilies can be expected to have the
369 same $\delta^{13}\text{C}$ values (van Hardenbroek et al. 2012), at least not in every lake. In a study
370 examining the sediments of a Siberian thermokarst lake, van Hardenbroek et al.
371 (2013) found that variations in $\delta^{13}\text{C}$ of *Chironomus*, Chironomini, Tanytarsini, and
372 *Daphnia* did not follow the same pattern as $\delta^{13}\text{C}$ of SOM. Only the chironomid
373 subfamily Orthocladiinae was characterized by $\delta^{13}\text{C}$ values very similar to those of
374 SOM. This could be explained by a preference of many Orthocladiinae for a diet of
375 periphyton or SOM, whereas the diet of *Chironomus*, Chironomini, Tanytarsini, and

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

376 *Daphnia* consisted, at least in part, of other components including planktonic algae
377 and bacteria. Van Hardenbroek et al. (2012; 2013) suggested that varying proportions
378 of methane oxidizing bacteria (MOB) in the diet of *Chironomus*, Chironomini,
379 Tanytarsini, and *Daphnia* may be at least partly responsible for the difference in $\delta^{13}\text{C}$
380 between the different invertebrate groups examined in their study. Furthermore,
381 Wooller et al. (2012) argued that methane-derived carbon led to lower $\delta^{13}\text{C}$ values of
382 chironomids and *Daphnia* in sections of a sediment record from arctic Alaska.
383 Frossard et al. (2014) also suggested that similar decreases of $\delta^{13}\text{C}$ values in two
384 Chironomini taxa, Tanypodinae, and pelagic Cladocera in a sediment record Lake
385 Annecy (France) were linked to increased availability of ^{13}C -depleted methane-
386 derived organic carbon. Chironomid larvae feeding partly on MOB can be strongly
387 ^{13}C -depleted, as shown for several species (Jones et al. 2008; Kiyashko et al. 2001;
388 Zemskaya et al. 2012).

389 For Strandsjön, relatively high $\delta^{13}\text{C}$ values for all chironomid taxa and for
390 *Daphnia* suggest that MOB may have been a less relevant food source than in the
391 lakes studied by Wooller et al. (2012), van Hardenbroek et al. (2013), and Frossard et
392 al. (2014). However, the finding that *Chironomus* and *C. mucedo* are relatively ^{13}C -
393 depleted could be explained by a higher importance of MOB in their diet than in the
394 food ingested by the other invertebrate groups. Larvae of several *Chironomus* species
395 have been shown to incorporate carbon originating from methane into their body
396 tissue (Jones et al. 2008). The feeding ecology of *C. mucedo* is still poorly studied,
397 but as a filter-feeder that can ingest particles in the μm range (Kaminski 1984), the
398 species may be able to feed on MOB in the water column. Similarly, the lower $\delta^{13}\text{C}$
399 values of chironomid remains of the groups Chironomini, Tanypodinae, and
400 Tanytarsini around AD ~1890, ~1920, ~1950, and ~1990 could be explained by

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

401 increasing importance of MOB in their diet. However, variations in $\delta^{13}\text{C}$ of
402 chironomid remains in Strandsjön were muted and well within the range of values
403 expected for photosynthetically produced organic matter (France 1995a,b; Meyers and
404 Terranes 2001; Bade et al. 2006). Therefore, the observed changes in chironomid $\delta^{13}\text{C}$
405 might also be caused by changes in the relative contribution of isotopically different
406 carbon sources, including allochthonous inputs, remains of aquatic macrophytes,
407 benthic algae, and periphyton that all have higher $\delta^{13}\text{C}$ values than pelagic algae
408 (France 1995a,b; Meyers and Terranes 2001). Changes in allochthonous inputs are not
409 obvious from the low and relatively stable C:N_{atomic} ratio of SOM that suggests
410 mainly autochthonous carbon sources. Benthic algae are not considered as an
411 important food source, as limited light penetration inhibits their growth in Strandsjön
412 with dissolved organic carbon concentrations as high as 20.8 mg L⁻¹ (van
413 Hardenbroek et al. 2012). Finally, it is unlikely that chironomid $\delta^{13}\text{C}$ responded to
414 changes in concentration or $\delta^{13}\text{C}$ of dissolved inorganic carbon in the water column,
415 as this would lead to changes in $\delta^{13}\text{C}$ of pelagic algae as well, and would be expected
416 to be registered also in the $\delta^{13}\text{C}$ of the filter-feeding *Daphnia* and bryozoans.
417 Remaining explanations for the observed variations in chironomid $\delta^{13}\text{C}$ include
418 changes in the biomass of macrophytes, periphyton, or MOB in Strandsjön which may
419 have affected the importance of these carbon sources in the diet of chironomid larvae.
420 $\delta^{13}\text{C}$ values of the filter-feeding *Daphnia*, *C. mucedo*, and *Plumatella* were
421 relatively constant in the older part of the record compared with chironomids; the
422 filter-feeders had a decreasing trend in $\delta^{13}\text{C}$ in the youngest sediments, followed by a
423 slight reversal to higher values in the uppermost samples. This pattern could be
424 related to changes in $\delta^{13}\text{C}$ of phytoplankton in the water column, since planktonic
425 algae can be expected to be an important component of the diet of *Daphnia* and

Bryozoa. The observed variations would be easiest to explain by changes in lake productivity, since $\delta^{13}\text{C}$ of phytoplankton in productive meso- to eutrophic lakes is often higher than $\delta^{13}\text{C}$ of phytoplankton growing in oligotrophic conditions (Brenner et al. 1999; Hollander et al. 1993). In Strandsjön, $\delta^{13}\text{C}$ of *Daphnia* and *C. mucedo* showed a maximum at ~AD 1960. $\delta^{13}\text{C}$ of *Plumatella* also showed relatively high values in this section of the record, although the maximum was not as pronounced as for the other two taxa and *Plumatella* was already characterized by relatively high $\delta^{13}\text{C}$ values in the period preceding ~AD 1960. A peak in lake nutrient concentration during this period agrees with what is known of the trophic state history of the lake: An increase in nutrient input as agriculture developed around the lake led to eutrophication in the 20th century, causing increasingly frequent algal blooms until the installation of a water treatment plant for the inflow in AD 1960, after which the water quality improved (Brunberg and Blomqvist 1998). In contrast to $\delta^{13}\text{C}$ of *Daphnia* and bryozoans, $\delta^{13}\text{C}$ of SOM did not seem to track the eutrophication history of the lake. This suggested that carbon isotopic analyses of invertebrate remains might be considerably more sensitive to past changes in $\delta^{13}\text{C}$ of organic matter produced in the water column than bulk SOM $\delta^{13}\text{C}$, at least in relatively small, shallow, and productive lakes such as Strandsjön.

Conclusions

Our results demonstrate that taxon-specific $\delta^{13}\text{C}$ records can be developed for a range of chitinous invertebrate remains found in lake sediments. In Strandsjön, our approach revealed a differential response of $\delta^{13}\text{C}$ of benthic taxa (chironomids and rhabdocoels) to past environmental change compared with $\delta^{13}\text{C}$ of taxa feeding predominantly on

suspended particulate organic matter (*Daphnia* and bryozoans). Hence, our study indicates that $\delta^{13}\text{C}$ of chitinous invertebrate remains can provide valuable additional information to $\delta^{13}\text{C}$ of SOM, as the latter cannot distinguish between carbon sources in different compartments of a lake. Additional studies relating the stable isotope composition of various aquatic invertebrates to the values of their fossilizing structures, as well as studies establishing the link between $\delta^{13}\text{C}$ of the diet and $\delta^{13}\text{C}$ of freshwater invertebrate tissues are urgently needed to improve the interpretation of invertebrate $\delta^{13}\text{C}$ in sediment records. More studies examining $\delta^{13}\text{C}$ of different organic invertebrate remains from the same sediment record are necessary to confirm that $\delta^{13}\text{C}$ of phytoplankton can be reliably traced by examining the carbon isotopic composition of the remains of filter-feeding *Daphnia* and Bryozoa, and that such analyses can reveal a different signal than the one captured by SOM or by the remains of zoobenthos. The results from Strandsjön demonstrate the potential of taxon-specific stable isotope analysis of invertebrate remains for environmental reconstructions and provide, to our knowledge, the taxonomically most diverse such study available to date. It is likely that similar taxon-specific records will soon be produced for stable isotopes of other elements (H, O, N, S) that can give further insights in the functioning of aquatic ecosystems over long time scales.

Acknowledgements

We thank Lotta Frisk Hagström and Kristina Eriksson for valuable historical information about the lake and Arndt Schimmelmann and three anonymous reviewers for their helpful suggestions to improve this manuscript. This research was supported by the Darwin Centre for Biogeosciences, the European Research Council (ERC)

476 Starting Grant project RECONMET (Project nr. 239858), and the Swedish Research
477 Council (Project no. VR 2006-3256).

478

479 **References**

480

481 Appleby PG (2001) Chronostratigraphic techniques in recent sediments. In: WM Last,
482 JP Smol (eds.), Tracking environmental change using lake sediments. Volume 1:
483 Basin analysis, coring, and chronological techniques. Kluwer Academic Publishers,
484 Dordrecht, pp. 171-203

485 Appleby PG, Oldfield F (1978) The calculation of lead-210 dates assuming a constant
486 rate of supply of unsupported ^{210}Pb to the sediment. *Catena* 5: 1-8

487 Bade D, Pace M, Cole J, Carpenter S (2006) Can algal photosynthetic inorganic
488 carbon isotope fractionation be predicted in lakes using existing models? *Aquat Sci*
489 68: 142-153

490 Barker PA, Hurrell ER, Leng MJ, Verschuren D, Conley D, Plessens B (2013) The
491 changing carbon cycle of lakes revealed by the stable isotopes of diatom silica: a 25
492 ka record from Lake Challa, Kilimanjaro. *Quaternary Sci Rev* 66: 55-63

493 Brenner M, Whitmore TJ, Curtis JH, Hodell DA, Schelske CL (1999) Stable isotope
494 ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) signatures of sedimented organic matter as indicators of historic lake
495 trophic state. *J Paleolimnol* 22: 205-221

496 Brooks SJ, Langdon PG, Heiri O (2007) The identification and use of Palaeartic
497 Chironomidae larvae in palaeoecology. QRA Technical Guide No. 10. Quaternary
498 Research Association, London

499 Brunberg A-K, Blomqvist P (1998) Vatten i Uppsala län 1997. Beskrivning,
500 utvärdering, åtgärdsförslag. Rapport nr 8/1998. Upplandsstiftelsen. [in Swedish]

501 DeNiro MJ, Epstein S (1978) Influence of diet on the distribution of carbon isotopes
502 in animals. *Geochim Cosmochim Acta* 42: 495-506

503 France RL (1995a) Carbon-13 enrichment in benthic compared to planktonic algae:
504 foodweb implications. *Mar Ecol Prog Ser* 124: 307-312

505 France RL (1995b) Stable isotopic survey of the role of macrophytes in the carbon
506 flow of aquatic foodwebs. *Vegetatio* 124: 67-72

507 Frey DG (1964) Remains of animals in Quaternary lake and bog sediments and their
508 interpretation. *Advances in Limnology* 2: 1-114

509 Frossard V, Belle S, Verneaux Vr, Millet L, Magny M (2013) A study of the $\delta^{13}\text{C}$
510 offset between chironomid larvae and their exuvial head capsules: implications for
511 palaeoecology. *J Paleolimnol* 50: 379-386

512

513 Frossard V, Verneaux V, Millet L, Jenny J-P, Arnaud F, Magny M, Perga M-E (2014)
514 Reconstructing long-term changes (150 years) in the carbon cycle of a clear-water
515 lake based on the stable carbon isotope composition ($\delta^{13}\text{C}$) of chironomid and
516 cladoceran subfossil remains. *Freshwat Biol* doi: 10.1111/fw.12304

517

518 Geller W, Müller H (1981) The filtration apparatus of Cladocera: Filter mesh-sizes
519 and their implications on food selectivity. *Oecologia* 49: 316-321

520 Heiri O, Schilder J, van Hardenbroek M (2012) Stable isotopic analysis of fossil
521 chironomids as an approach to environmental reconstruction: State of development
522 and future challenges. *Proceedings of the 18th International Symposium on*
523 *Chironomidae*, Trondheim 4-6. August 2011, *Fauna Norvegica* 31: 7-18

524 Herzsich U, Mischke S, Meyer H, Plessen B, Zhang C (2010) Lake nutrient
525 variability inferred from elemental (C, N, S) and isotopic ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$) analyses of
526 aquatic plant macrofossils. *Quaternary Sci Rev* 29: 2161-2172

527 Hollander DJ, McKenzie JA, Hsu KJ, Huc AY (1993) Application of an eutrophic
528 lake model to the origin of ancient organic-carbon-rich sediments. *Global*
529 *Biogeochem Cycles* 7: 157-179

530 Hurrell ER, Barker PA, Leng MJ, Vane CH, Wynn P, Kendrick CP, Verschuren D,
531 Street-Perrott FA (2011) Developing a methodology for carbon isotope analysis of
532 lacustrine diatoms. *Rapid Commun Mass Sp* 25: 1567-1574

533 Jennings JB (1957) Studies on feeding, digestion, and food storage in free-living
534 flatworms (Platyhelminthes: Turbellaria). *Biological Bulletin* 112: 63-80

535 Jones RI, Carter CE, Kelly A, Ward S, Kelly DJ, Grey J (2008) Widespread
536 contribution of methane-cycle bacteria to the diets of lake profundal chironomid
537 larvae. *Ecology* 89: 857-864

538 Kaminski M (1984) Food composition of three bryozoan species (Bryozoa,
539 Phylactolaemata) in a mesotrophic lake. *Polish Archive of Hydrobiology* 31: 45-53

540 Kiyashko SI, Narita T, Wada E (2001) Contribution of methanotrophs to freshwater
541 macroinvertebrates: evidence from stable isotope ratios. *Aquat Microb Ecol* 24: 203-
542 207

543 Kolasa J, Tyler S (2010) Flatworms: Turbellaria and Nemertea. In: JH Thorp, AP
544 Covich (eds.), *Ecology and classification of North American freshwater invertebrates*.
545 Academic Press, London, pp. 143-161

546 Leavitt PR, Brock CS, Ebel C, Patoine A (2006) Landscape-scale effects of urban
547 nitrogen on a chain of freshwater lakes in central North America. *Limnol Oceanogr*
548 51: 2262-2277

549 Legendre P, Legendre L (1998) *Numerical ecology*. Developments in environmental
550 modelling 20. Elsevier Science, Amsterdam, 853 pp

551

- 552 Leng MJ, Lamb AL, Marshall JD, Wolfe BB, Jones MD, Holmes JA, Arrowsmith C
553 (2005) Isotopes in lake sediments. In: MJ Leng (ed.), Isotopes in palaeoenvironmental
554 research. Springer, Dordrecht, pp. 147-184
- 555 Luther A (1955) Die Dalyelliiden (Turbellaria Neorhabdocoela), eine monografie.
556 von Tilgmann, Helsingfors, 337 pp
- 557 Macko SA, Helleur R, Hartley G, Jackman P (1990) Diagenesis of organic matter - a
558 study using stable isotopes of individual carbohydrates. Org Geochem 16: 1129-1137
559
- 560 Merritt RW, Cummins KW, Berg MB (2008) An introduction to the Aquatic Insects
561 of North America. Kendall and Hunt, Dubuque, 1158 pp
- 562 Meyers PA, Lallier-Vergès E (1999) Lacustrine sedimentary organic matter records of
563 Late Quaternary paleoclimates. J Paleolimnol 21: 345-372
- 564 Meyers PA, Teranes JL (2001) Sediment Organic Matter. In: WM Last, JP Smol
565 (eds.), Tracking environmental change using lake sediments. Volume 2: Physical and
566 Geochemical Techniques. Kluwer Academic Publishers, Dordrecht, pp. 239-269
- 567 Mihuc T, Toetz D (1994) Determination of Diets of Alpine Aquatic Insects Using
568 Stable Isotopes and Gut Analysis. The American Midland Naturalist 131: 146-155
- 569 Moller Pillot HKM (2009) Chironomidae Larvae, Biology and ecology of the
570 Chironomini. KNNV Publishing, Zeist, 272 pp
- 571 Moog O (2002) Fauna Aquatica Austriaca. Wasserwirtschaftskataster,
572 Bundesministerium für Land- und Forstwirtschaft, Umwelt und Wasserwirtschaft,
573 Vienna
- 574 Okamura B, Hatton-Ellis T (1995) Population biology of bryozoans: correlates of
575 sessile, colonial life histories in freshwater habitats. Cell Mol Life Sci 51: 510-525
- 576 Perga M-E (2010) Potential of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of cladoceran subfossil exoskeletons for
577 paleo-ecological studies. J Paleolimnol 44: 387-395
- 578 Perga M-E (2011) Taphonomic and early diagenetic effects on the C and N stable
579 isotope composition of cladoceran remains: implications for paleoecological studies. J
580 Paleolimnol 46: 203-213
- 581 Rau GH (1980) Carbon-13/Carbon-12 Variation in Subalpine Lake Aquatic Insects:
582 Food Source Implications. Can J Fish Aquat Sci 37: 742-746
- 583 Schimmelmann A (2011) Carbon, Nitrogen and Oxygen Stable Isotope Ratios in
584 Chitin. In: NS Gupta (ed.), Chitin. Springer Netherlands, pp. 81-103
- 585 Stevenson FJ (1982) Extraction, fractionation, and general chemical composition of
586 soil organic matter. In: FJ Stevenson (ed.), Humus Chemistry. Genesis, Composition,
587 Reactions, John Wiley and Sons, New York, pp. 26-54
- 588 Vallenduuk H, Moller Pillot HKM (2007) Chironomidae larvae, General introduction
589 and Tanypodinae. KNNV Publishing, Zeist, 172 pp

- 590 Vanderkerkhove J, Declerck S, Vanhove M, Brendonck L, Jeppesen E, Porcuna JM,
591 De Meester L (2004) Use of ehippial morphology to assess richness of anomopods:
592 potentials and pitfalls. *J Limnol* 63: 75-84
- 593 Vander Zanden MJ, Rasmussen JB (1999) Primary consumer $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ and the
594 trophic position of aquatic consumers. *Ecology* 80: 1395-1404
- 595 van Hardenbroek M, Heiri O, Grey J, Bodelier P, Verbruggen F, Lotter A (2010a)
596 Fossil chironomid $\delta^{13}\text{C}$ as a proxy for past methanogenic contribution to benthic food
597 webs in lakes? *J Paleolimnol* 43: 235-245
- 598 van Hardenbroek M, Heiri O, Lotter A (2010b) Efficiency of different mesh sizes for
599 isolating fossil chironomids for stable isotope and radiocarbon analyses. *J Paleolimnol*
600 44: 721-729
- 601 van Hardenbroek M, Lotter AF, Bastviken D, Duc NT, Heiri O (2012) Relationship
602 between $\delta^{13}\text{C}$ of chironomid remains and methane flux in Swedish lakes. *Freshwat*
603 *Biol* 57: 166-177
- 604 van Hardenbroek M, Heiri O, Parmentier FJW, Bastviken D, Ilyashuk BP, Wiklund
605 JA, Hall RI, Lotter AF (2013) Evidence for past variations in methane availability in a
606 Siberian thermokarst lake based on $\delta^{13}\text{C}$ of chitinous invertebrate remains. *Quaternary*
607 *Sci Rev* 66: 74-84
- 608 Verbruggen F, Heiri O, Reichart GJ, De Leeuw J, Nierop K, Lotter A (2010) Effects
609 of chemical pretreatments on $\delta^{18}\text{O}$ measurements, chemical composition, and
610 morphology of chironomid head capsules. *J Paleolimnol* 43: 857-872
- 611 Wood TS, Okamura B (2005) A new key to freshwater bryozoans of Britain, Ireland
612 and continental Europe, with notes on their ecology. *Freshwater Biological*
613 *Association, London*, 113 pp
- 614 Wooller M, Pohlman J, Gaglioti B, Langdon P, Jones M, Walter Anthony K, Becker
615 K, Hinrichs K-U, Elvert M (2012) Reconstruction of past methane availability in an
616 Arctic Alaska wetland indicates climate influenced methane release during the past
617 ~12,000 years. *J Paleolimnol* 48: 27-42
- 618 Wooller MJ, Wang Y, Axford Y (2008) A multiple stable isotope record of Late
619 Quaternary limnological changes and chironomid paleoecology from northeastern
620 Iceland. *J Paleolimnol* 40: 63-77
- 621
622 Zenskaya T, Sitnikova T, Kiyashko S, Kalmychkov G, Pogodaeva T, Mekhanikova I,
623 Naumova T, Shubenkova O, Chernitsina S, Kotsar O, Chernyaev E, Khlystov O
624 (2012) Faunal communities at sites of gas- and oil-bearing fluids in Lake Baikal. *Geo-*
625 *Mar Lett* 32: 437-451

Figure captions

Fig. 1 Site location and bathymetric map of Strandsjön (redrawn after Brunberg and Blomqvist 1998). Coring site is indicated by X, inlets and outlets are indicated by arrows.

Fig. 2 Activity of ^{210}Pb (left) and ^{137}Cs (middle) and the age model (right) for the sediment record from Strandsjön based on a constant rate of supply model for ^{210}Pb with the 4.5 cm peak in ^{137}Cs activity as reference point for the 1986 Chernobyl accident.

Fig. 3 $\delta^{13}\text{C}$ values of invertebrate remains and geochemistry of sedimentary organic matter ($\delta^{13}\text{C}$, organic carbon content, and $\text{C:N}_{\text{atomic}}$ ratio) in the sediment record from Strandsjön. Invertebrate taxa are indicated by open triangles (*Chironomus*), open circles (Chironomini), open grey squares (Tanytarsini), open diamonds (Tanypodinae), solid circles (*Daphnia*), solid triangles (*Cristatella mucedo*), solid diamonds (*Plumatella*), and solid grey squares (Rhabdocoela).

Fig. 4 Boxplots with $\delta^{13}\text{C}$ values of the different invertebrate taxa in this study and of sedimentary organic matter (SOM).

Figure 1
[Click here to download Figure: Fig 1 Location.eps](#)




Figure 2
[Click here to download Figure: Fig 2 Age depth v2.eps](#)




Figure 4
[Click here to download Figure: Fig 4 boxplot v2.eps](#)





Figure 3
[Click here to download Figure: Fig 3 Downcore Strandsjon v5.eps](#)



Supplementary table 1: $\delta^{13}\text{C}$, number of remains, and weight (in microgram) for each invertebrate sample. $\delta^{13}\text{C}$, $\delta^{15}\text{N}$, $\%C_{\text{weight}}$, $\%N_{\text{weight}}$ and $C:N_{\text{atomic}}$ ratio for sedimentary organic matter (SOM)

Depth (cm)	<i>Chironomus</i>			Chironomini			Tanytarsini			Tanypodinae			<i>Daphnia</i>			<i>Cristatella mucedo</i>			<i>Plumatella</i>			Rhabdocoela			SOM				
	$\delta^{13}\text{C}$	no.	mass	$\delta^{13}\text{C}$	no.	mass	$\delta^{13}\text{C}$	no.	mass	$\delta^{13}\text{C}$	no.	mass	$\delta^{13}\text{C}$	no.	mass	$\delta^{13}\text{C}$	no.	mass	$\delta^{13}\text{C}$	no.	mass	$\delta^{13}\text{C}$	no.	mass	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	$\%C_{\text{weight}}$	$\%N_{\text{weight}}$	$C:N_{\text{atomic}}$
0-2				-30.53	21	18	-30.3	39	17	-31.9	15	10	-31.45	13	15	-33.7	1.25	29	-31.27	30	17	-29.1	40	36	-32.8	3.4	12.6	1.4	10.6
2-3	-31.7	19	18	-32.2	41	29	-31.2	41	20	-31.9	19	9	-32.8	31	28				-32.1	40	21	-29.2	40	37	-32.8	3.4	12.3	1.3	10.7
4-5	-33.2	27	24	-32.1	41	23	-31.9	47	15	-32.4	23	15	-32.3	32	29	-34.2	1	22	-33.1	40	29	-29.2	40	33	-32.5	3.4	12.2	1.3	10.7
6-7	-33.0	31	36	-31.4	33	22	-31.8	45	22	-31.8	22	15	-31.7	36	42	-35.1	1.25	43	-30.7	40	24	-30.0	40	40	-31.4	3.5	11.1	1.2	11.0
8-9	-32.6	31	41	-31.4	38	40	-31.5	45	28				-31.4	41	36				-30.8	45	21	-29.2	40	47	-31.2	3.5	10.5	1.1	11.1
10-11	-31.7	12	26	-31.6	31	29	-32.2	45	28	-32.2	31	27	-30.5	40	37	-31.3	1.5	59	-29.1	40	23	-29.2	40	48	-31.0	3.3	10.9	1.1	11.5
12-13	-32.6	8	11	-33.4	44	39	-33.9	47	32	-33.8	19	22	-30.5	41	34	-32.6	1.25	21	-29.2	40	24	-30.2	40	56	-31.5	2.8	13.6	1.3	12.0
14-15	-34.6	12	19	-32.6	16	12	-33.5	47	23	-32.8	20	17	-30.9	40	29	-31.4	1.5	34	-28.2	40	18	-29.7	40	48	-31.4	2.6	15.3	1.5	12.2
16-17				-32.0	38	38	-32.5	35	16				-31.3	30	44				-29.6	40	16	-29.6	40	47					
18-19	-34.2	15	24	-33.2	38	36	-32.7	50	21	-32.9	25	21	-31.2	41	54				-28.9	45	18	-29.7	40	53	-30.8	2.4	15.9	1.5	12.4
20-21				-33.5	39	42	-32.1	45	25	-33.4	33	27	-31.6	31	49	-33.5	2	34	-28.9	35	15	-30.0	40	51	-30.6	2.7	15.0	1.4	13.4
22-23				-32.4	35	42	-32.5	46	27	-31.7	14	15	-32.1	32	48	-32.6	1.25	22	-29.5	40	15	-29.7	40	61	-30.6	2.7	15.8	1.5	12.0
24-25				-31.9	39	42	-32.3	45	23	-31.2	15	20	-31.9	29	40				-29.1	40	13	-29.5	40	53	-30.4	2.6	16.3	1.5	12.0
26-27				-31.8	40	49	-33.5	45	29	-32.0	19	27	-32.1	30	39	-33.1	1	20	-28.2	42	15	-29.3	40	53	-30.5	2.6	16.6	1.6	11.9
28-29				-32.7	38	30	-33.8	45	24	-32.6	26	33	-32.0	31	48				-29.4	40	17	-29.1	40	46	-30.5	2.4	18.9	1.9	10.4
30-32				-31.2	38	28	-32.3	45	21	-32.0	24	30	-32.0	31	42	-31.6	2	17	-28.6	40	13	-29.3	40	53	-30.1	2.4	20.0	1.9	11.7
32-33	-30.8	8	9	-31.1	31	32	-32.0	45	25	-31.0	18	26	-32.2	30	39	-31.2	0.5	10	-28.1	40	14	-29.0	40	44	-30.0	2.3	20.1	1.8	12.7
34-35				-31.3	37	23	-32.3	45	24	-31.5	25	32	-32.0	30	34	-34.1	1	21	-28.0	42	16	-28.4	40	38	-30.3	2.3	19.1	1.8	12.9

Below recommended weight of 20 microgram